

Table 17: Presence of Genetically Modified Maize (2009/2010)

REGION	RSA Grade	MON810, % (LOD: 0.15%)	NK603 (Roundup Ready) % (LOD: 0.25%)	REGION	RSA Grade	MON810, % (LOD: 0.15%)	NK603 (Roundup Ready) % (LOD: 0.25%)
10	YM1	>2	>1.8	25	WM1	>2	<0.25
11	WM1	>2	0.4	25	WM2	>2	<0.25
12	WM1	>2	<0.25	25	WM2	>2	1.3
12	WM3	>2	0.8	25	YM1	>2	0.3
13	WM3	>2	0.6	25	YM1	>2	0.4
13	YM1	>2	1.7	26	WM1	>2	<0.25
14	WM2	>2	<0.25	26	WM2	>2	0.4
14	WM2	>2	0.8	26	WM3	>2	<0.25
14	WM2	>2	1.2	26	YM1	>2	>1.8
14	YM1	>2	0.6	26	YM1	>2	<0.25
15	WM2	>2	<0.25	27	WM2	>2	<0.25
15	YM1	>2	>1.8	27	WM2	>2	1.2
16	WM1	>2	0.3	27	YM2	>2	<0.25
16	WM1	>2	<0.25	28	WM1	>2	>1.8
16	YM2	>2	1.3	28	WM1	>2	>1.8
17	WM1	>2	0.6	28	WM1	>2	1.5
17	WM2	>2	<0.25	28	YM1	>2	>1.8
17	WM2	>2	0.9	28	YM1	>2	<0.25
17	WM2	>2	1.8	28	YM1	>2	>1.8
18	WM1	>2	>1.8	28	YM2	1.0	<0.25
18	WM1	>2	0.8	28	YM2	>2	<0.25
18	WM2	>2	<0.25	28	YM3	>2	0.4
18	YM1	>2	1.2	29	WM1	>2	0.5
18	YM1	>2	0.5	29	WM1	>2	<0.25
19	WM2	>2	<0.25	29	WM1	>2	1.7
19	YM2	>2	1.1	29	WM2	>2	<0.25
20	WM2	>2	1.2	29	YM1	>2	<0.25
20	WM3	>2	1.4	29	YM1	0.3	<0.25
21	WM1	>2	>1.8	30	WM1	>2	>1.8
21	WM1	>2	>1.8	30	WM1	>2	<0.25
21	WM1	>2	0.7	30	WM1	<0.15	<0.25
21	WM1	>2	1.0	31	WM1	<0.15	<0.25
22	WM1	>2	1.4	31	YM2	>2	>1.8
22	WM2	>2	1.4	31	YM2	1.8	<0.25
22	WM2	>2	0.5	32	WM1	<0.15	<0.25
23	WM1	>2	0.3	32	WM1	>2	<0.25
23	WM3	>2	0.3	33	WM1	>2	0.4
23	YM1	>2	<0.25	34	WM1	>2	<0.25
23	YM1	>2	<0.25	34	YM1	<0.15	<0.25
23	YM2	>2	>1.8	34	YM1	>2	0.7
24	WM1	>2	>1.8	35	WM1	>2	>1.8
24	WM1	>2	<0.25	35	YM1	>2	>1.8
24	WM1	>2	<0.25	36	WM1	0.5	>1.8
24	WM1	>2	0.8	36	WM1	>2	<0.25
24	WM1	>2	<0.25	36	YM1	>2	0.8
n	Season	% Samples positive for MON810 (Bt)		n	Season	% Samples positive for NK603 (RUR)	
90	2009/2010	96		90	2009/2010	61	
90	2008/2009	91		90	2008/2009	90	
100	2007/2008	95		100	2007/2008	69	
90	2006/2007	97		90	2006/2007	59	
90	2005/2006	91		90	2005/2006	31	
100	2004/2005	78		100	2004/2005	31	