

BREAD WHEAT GRADING TABLE 2009/2010

Grade	Minimum			Maximum percentage permissible deviation (m/m)									
				A	B	C	D	E	F	G	H	I	J
	Hectolitre mass, kg	Falling number, seconds	Protein content, %	Heavily frost damaged kernels	Field fungi	Storage fungi	Screenings	Other grain and unthreshed ears	Gravel, stones, turf and glass	Foreign matter plus F	Heat damaged kernels	Damaged kernels plus H	Combined deviations (D+E+G+I)
Grade 1	77	220	12	5	2	0.5	3	1	0.5	1	0.5	2	5
Grade 2	76	220	11	5	2	0.5	3	1	0.5	1	0.5	2	5
Grade 3	74	220	10	5	2	0.5	3	1	0.5	1	0.5	2	5
Grade 4	72	200	9	5	2	0.5	3	1	0.5	1	0.5	2	5
Utility grade	70	150	8	10	2	0.5	10	4	0.5	3	0.5	5	10
Other Wheat	<70	<150	<8	>10	>2	>0.5	>10	>4	>0.5	>3	>0.5	>5	>10
Minimum size of working samples	1 kg	300 g clean	Apparatus instructions	25 g sifted	25 g sifted	25 g sifted	500 g unsifted	50 g sifted	100 g sifted	100 g sifted	100 g sifted	25 g sifted	-

MYCOTOXIN RESULTS FOR THE 2009/2010 SEASON

Region	Class and Grade	Aflatoxin	Deoxynivalenol	Ochratoxin A
		ppb LOD < 2.0	ppm LOD < 0.25	ppb LOD < 1.0
1	B3	0	0	0
2	B2	2	0	1
3	B3	0	0	1
4	B4	0	0	0
5	B3	0	0	0
6	B1	0	0	1
7	B1	2	0	0
10	B1	0	0	0
11	B1	0	0	0
12	UT	0	0	0
14	B2	2	0	0
15	B2	0	0	0
16	B1	0	0.27	0
17	B1	0	0	0
18	B2	0	0	0
19	B2	0	0.48	0
20	B3	2	0	0
21	B4	4	0	0
22	B1	2	0	0
23	B4	0	0	0
24	B1	0	0	0
25	B2	2	0	0
26	B1	2	0	0
27	B3	0	0	0
28	B1	2	0	0
32	B2	3	0	1
33	B2	2	0.42	0
34	B1	0	0.48	1
35	B1	2	0	0
36	B1	3	0	0
Average 2009/2010 [max. value]		1.00 [4]	0.05 [0.48]	0.17 [1]
Average 2008/2009 [max. value]		1.23 [3]	0.47 [3]	0.03 [1]
Average 2007/2008 [max. value]		0.33 [5]	1.36 [2.7]	0.33 [2.8]

Please note:

All results <LOD and non detected are reported as 0.
LOD: Limit of detection, see table.