




**TABLE 14: COMPARISON OF THE AVERAGE NUTRITIONAL VALUES BETWEEN WHITE AND YELLOW MAIZE OVER THE PAST TEN SEASONS**

Season	Number of samples	White maize			Number of samples	Yellow maize		
		Fat	Protein	Starch		Fat	Protein	Starch
1998/99	256	4.1	9.2	71.8	189	4.2	9.5	72.1
1999/00	493	4.0	8.1	71.9	407	4.1	8.0	72.0
2000/01	522	4.2	8.8	70.2	378	4.2	8.7	70.5
2001/02	471	4.2	8.9	71.4	429	4.1	8.9	71.7
2002/03	517	4.1	9.2	71.4	383	4.1	9.2	72.0
2003/04	599	4.0	9.1	71.2	301	4.0	9.0	71.1
2004/05	601	4.0	8.9	71.1	399	3.8	8.6	71.7
2005/06	593	4.0	8.5	71.1	307	3.9	8.4	71.5
2006/07	563	3.9	9.3	72.9	337	3.5	9.5	73.3
2007/08	483	4.0	8.6	71.9	417	3.6	8.4	72.3
<b>Weighted averages</b>		4.0	8.9	71.5		3.9	8.8	71.8

**TABLE 15: AVERAGE NUTRITIONAL VALUES OF SOUTH AFRICAN MAIZE (1998/99 - 2007/08)**

Season	Number of samples	Fat %	Protein %	Starch %
1998/99	445	4.2	9.3	71.9
1999/00	900	4.0	8.1	71.9
2000/01	900	4.2	8.8	70.3
2001/02	900	4.2	8.9	71.5
2002/03	900	4.1	9.2	71.6
2003/04	900	4.0	9.1	71.1
2004/05	1000	3.9	8.8	71.3
2005/06	900	4.0	8.4	71.2
2006/07	900	3.7	9.4	73.0
2007/08	900	3.8	8.5	72.1
<b>Weighted averages</b>		4.0	8.8	71.6



**Please note:** Different starch methods have been used over years and data have been corrected accordingly.